

MEDICATION GUIDE

Zolpimist Oral Spray (C-IV) (zolpidem tartrate) Spray, Metered for Oral Use

Read the Medication Guide that comes with Zolpimist before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about your medical condition or treatment.

What is the most important information I should know about Zolpimist?

After taking Zolpimist, you may get up out of bed while not being fully awake and do an activity that you do not know you are doing. The next morning, you may not remember that you did anything during the night. You have a higher chance for doing these activities if you drink alcohol or take other medicines that make you sleepy with Zolpimist. Reported activities include:

- driving a car ("sleep-driving")
- making and eating food
- talking on the phone
- having sex
- sleep-walking

Call your doctor right away if you find out that you have done any of the above activities after taking Zolpimist. Important:

1. Take Zolpimist exactly as prescribed

- Do not take more Zolpimist than prescribed.
- Take Zolpimist right before you get in bed, not sooner.

2. Do not take Zolpimist if you:

- Drink alcohol
- Take other medicines that can make you sleepy. Talk to your doctor about all of your medicines. Your doctor will tell you if you can take Zolpimist with your other medicines.
- Cannot get a full night sleep

What is Zolpimist?

Zolpimist is a sedative-hypnotic (sleep) medicine. Zolpimist is used in adults for the short-term treatment of a sleep problem called insomnia. Symptoms of insomnia include:

- Trouble falling asleep

Zolpimist is not for children.

Zolpimist is a federally controlled substance (C-IV) because it can be abused and lead to dependence. Keep Zolpimist in a safe place to prevent misuse and abuse. Selling or giving away Zolpimist may harm others, and is against the law. Tell your doctor if you have ever abused or have been dependent on alcohol, prescription medicines, or street drugs.

Who should not take Zolpimist? Do not take Zolpimist if you have had an allergic reaction to zolpidem (Ambien, Ambien CR, Zolpimist). Some signs of allergic reaction may be swelling of the face, a feeling of the throat closing, or difficulty breathing shortly after taking Zolpidem.

See the end of this Medication Guide for a complete list of ingredients in Zolpimist.

Zolpimist may not be right for you. Before starting Zolpimist, tell your doctor about all of your health conditions, including if you:

- have a history of depression, mental illness, or suicidal thoughts
- have a history of drug or alcohol abuse or addiction
- have kidney or liver disease
- have a lung disease or breathing problems
- are pregnant, planning to become pregnant, or breastfeeding

Tell your doctor about all the medicines you take including prescription and nonprescription medicines, vitamins, and herbal supplements. Medicines can interact with each other, sometimes causing serious side effects. **Do not take Zolpimist with other medicines that can make you sleepy.**

Know the medicines you take. Keep a list of your medicines with you to show your doctor and pharmacist each time you get a new medicine.

How should I take Zolpimist?

- Take Zolpimist exactly as prescribed. Do not take more Zolpimist than prescribed for you.
- **Take Zolpimist right before you get into bed.**
- **Do not take Zolpimist unless you are able to stay in bed a full night (7-8 hours) before you must be active again.**
- For faster sleep onset, Zolpimist should NOT be taken with or immediately after a meal.

- Call your doctor if your insomnia worsens or is not better within 7 to 10 days. This may mean that there is another condition causing your sleep problem.

- If you take too much Zolpimist or overdose, call your doctor or poison control center right away, or get emergency treatment.

What are the possible side effects of Zolpimist?

Serious side effects of Zolpimist include:

- **getting out of bed while not being fully awake and doing an activity that you do not know you are doing.** (See "What is the most important information I should know about Zolpimist?")

- **abnormal thoughts and behavior.** Symptoms include more outgoing or aggressive behavior than normal, confusion, agitation, hallucinations, worsening of depression, and suicidal thoughts or actions.

- **memory loss**

- **anxiety**

- **severe allergic reactions.** Symptoms include swelling of the tongue or throat, trouble breathing, and nausea and vomiting. Get emergency medical help if you get these symptoms after taking Zolpimist.

Call your doctor right away if you have any of the above side effects or any other side effects that worry you while using Zolpimist.

The most common side effects of Zolpimist are:

- drowsiness

- dizziness

- diarrhea

- "drugged feelings"

- You may still feel drowsy the next day after taking Zolpimist. **Do not drive or do other dangerous activities after taking Zolpimist until you feel fully awake.**

After you stop taking a sleep medicine, you may have symptoms for 1 or 2 days such as: tiredness, trouble sleeping, nausea, flushing, lightheadedness, uncontrolled crying, vomiting, stomach cramps, panic attack, nervousness, and stomach area pain.

These are not all the side effects of Zolpimist. Ask your doctor or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Zolpimist?

- Store Zolpimist in an upright position at 59 F to 86 F (15 °C to 30 °C).

- Do not freeze.

- Avoid prolonged product exposure above 86 F (30 °C).
- The child-resistant container should be thrown away when the 60 sprays have been used.

Keep Zolpimist and all medicines out of reach of children.

General information about Zolpimist

- Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

- Do not use Zolpimist for a condition for which it was not prescribed.

- Do not share Zolpimist with other people, even if you think they may have the same symptoms that you have. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about Zolpimist. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Zolpimist that is written for healthcare professionals. For more information about Zolpimist, call 1-800-527-1955.

What are the ingredients in Zolpimist?

Active Ingredient: Zolpidem tartrate

Inactive Ingredients: artificial cherry flavor, benzoic acid, citric acid monohydrate, hydrochloric acid, neotame, propylene glycol, and purified water.

Rx Only

Revised September, 2010

This Medication Guide has been approved by U.S. Food and Drug Administration.

ECR Pharmaceuticals Richmond, Virginia 23255